

STEIN'S METHOD, HEAT KERNEL, AND TRACES OF POWERS OF ELEMENTS OF COMPACT LIE GROUPS

JASON FULMAN

ABSTRACT. Combining Stein's method with heat kernel techniques, we show that the trace of the j th power of an element of $U(n, \mathbb{C})$, $USp(n, \mathbb{C})$ or $SO(n, \mathbb{R})$ has a normal limit with error term of order j/n . In contrast to previous works, here j may be growing with n . The technique should prove useful in the study of the value distribution of approximate eigenfunctions of Laplacians.

1. INTRODUCTION

There is a large literature on the traces of powers of random elements of compact Lie groups. One of the earliest results is due to Diaconis and Shahshahani [5]. Using the method of moments, they show that if M is random from the Haar measure of the unitary group $U(n, \mathbb{C})$, and $Z = X + iY$ is a standard complex normal with X and Y independent, mean 0 and variance $\frac{1}{2}$ normal variables, then for $j = 1, 2, \dots$, the traces $Tr(M^j)$ are independent and distributed as $\sqrt{j}Z$ asymptotically as $n \rightarrow \infty$. They give similar results for the orthogonal group $O(n, \mathbb{R})$ and the group of unitary symplectic matrices $USp(2n, \mathbb{C})$. The moment computations of [5] use representation theory. It is worth noting that there are other approaches to their moment computations: [19] uses a version of integration by parts, [11] uses the combinatorics of cumulant expansions, and [3] uses an extended Wick calculus. We mention that traces of powers of random matrices have been studied for other matrix ensembles too ([2],[7],[28]).

Concerning the error in the normal approximation, Diaconis conjectured that for fixed j , it decreases exponentially or even subexponentially in n . In an ingenious paper (which is quite technical and seems tricky to apply to other settings), Stein [30] uses an iterative version of "Stein's method" to show that for j fixed, $Tr(M^j)$ on $O(n, \mathbb{R})$ is asymptotically normal with error $O(n^{-r})$ for any fixed r . Johansson [12] proved Diaconis' conjecture for classical compact Lie groups using Toeplitz determinants and a very detailed analysis of characteristic functions. Duits and Johansson [6] allow j to grow with n in the unitary case, but do not obtain error terms. We also note that in the unitary case when $j \geq n$, the situation is not so interesting, since by work of Rains [21], the eigenvalues of M^j are simply n independent points from the unit circle (and he proves analogous results for other compact Lie groups).

Date: May 4, 2010.

Key words and phrases. Random matrix, Stein's method, heat kernel.

Dedicated to Thuy Le, on the occasion of our tenth anniversary.

The current paper studies the distribution of $Tr(M^j)$ using Stein's method and heat kernel techniques. This is a follow-up work to the paper [9], which used Stein's method and character theory to study the distribution of $\chi(M)$, where χ is the character of an irreducible representation; the functions $Tr(M^j)$ are not irreducible characters for $j > 1$, so do not fit into the framework of [9]. It should also be mentioned that the heat kernel is a truly remarkable tool appearing in many parts of mathematics (see the article [13] for a spirited defense of this statement with many references), and we suspect that the blending of heat kernel techniques with Stein's method will be useful for other problems.

In earlier work, Meckes [17], used Stein's method to study eigenfunctions of the Laplacian (a topic of interest in quantum chaos and arithmetic [27], among other places). We note two differences with her work. First, she uses geodesic flows and Liouville measure instead of heat kernels. Second, her infinitesimal version of Stein's method [17], [18] uses an exchangeable pair of random variables (W, W_ϵ) with the conditional expectation $\mathbb{E}[W_\epsilon - W|W]$ divided by ϵ^2 approximately proportional to W as $\epsilon \rightarrow 0$. In the current paper the natural condition is that $\mathbb{E}[W_\epsilon - W|W]$ divided by ϵ is approximately proportional to W as $\epsilon \rightarrow 0$.

We do use some moment computations from [5], but as is typical with Stein's method, only a few low order moments are needed. It should also be mentioned that the constants in our error terms can be made completely explicit (for instance in the unitary case we prove a bound of $\frac{22j}{n}$), but we do not work out the other constants as the bookkeeping is tedious and the true convergence rate is likely to be of a sharper order. As to future work, we note that more general linear combinations of traces of powers do satisfy central limit theorems (see [4], [12], [29] for precise conditions); obtaining good error terms by our techniques (or other methods) may be quite tricky and is an important problem.

The organization of this paper is as follows. Section 2 gives background on both Stein's method and the heat kernel. Section 3 treats the orthogonal groups, Section 4 treats the symplectic groups, and Section 5 treats the unitary groups.

2. STEIN'S METHOD AND THE HEAT KERNEL

In this section we briefly review Stein's method for normal approximation, using the method of exchangeable pairs [31]. One can also use couplings to prove normal approximations by Stein's method (see [22] for a survey), but the exchangeable pairs approach is effective for our purposes. For a survey discussing both exchangeable pairs and couplings, the paper [23] can be consulted.

Two random variables W, W' on a state space X are called exchangeable if the distribution of (W, W') is the same as the distribution of (W', W) . As is typical in probability theory, let $\mathbb{E}(A|B)$ denote the expected value of A given B . The following result of Rinott and Rotar [24] uses an exchangeable pair (W, W') to prove a central limit theorem for W .

Theorem 2.1. ([24]) *Let (W, W') be an exchangeable pair of real random variables such that $\mathbb{E}(W) = 0, \mathbb{E}(W^2) = 1$ and $\mathbb{E}(W'|W) = (1-a)W + R(W)$ with $0 < a < 1$. Then for all real x_0 ,*

$$\begin{aligned} & \left| \mathbb{P}(W \leq x_0) - \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{x_0} e^{-\frac{x^2}{2}} dx \right| \\ & \leq \frac{6}{a} \sqrt{\text{Var}(\mathbb{E}[(W' - W)^2|W])} + 19 \frac{\sqrt{\mathbb{E}(R^2)}}{a} + 6 \sqrt{\frac{1}{a} \mathbb{E}|W' - W|^3}. \end{aligned}$$

In practice, it can be quite challenging to construct exchangeable pairs satisfying the hypotheses of Theorem 2.1, and such that the error terms are tractable and small.

Lemma 2.2 is a known inequality (already used in the monograph [31]) and useful because often the right hand sides are easier to compute or bound than the left hand sides. We include the short proof. Here M is an element of the state space X (in this paper X is a compact Lie group and M a matrix in X).

Lemma 2.2. (1) $\text{Var}(\mathbb{E}[(W' - W)^2|W]) \leq \text{Var}(\mathbb{E}[(W' - W)^2|M])$.
 (2) *With notation as in Theorem 2.1, letting $\mathbb{E}(W'|M) = (1-a)W + R(M)$, one has that $\mathbb{E}(R(W)^2) \leq \mathbb{E}(R(M)^2)$.*

Proof. Jensen's inequality states that if g is a convex function, and Z a random variable, then $g(\mathbb{E}(Z)) \leq \mathbb{E}(g(Z))$. There is also a conditional version of Jensen's inequality (Section 4.1 of [8]) which states that for any σ subalgebra F of the σ -algebra of all subsets of X ,

$$\mathbb{E}(g(\mathbb{E}(Z|F))) \leq \mathbb{E}(g(Z)).$$

Part 1 now follows by setting $g(t) = t^2$, $Z = \mathbb{E}((W' - W)^2|M)$, and letting F be the σ -algebra generated by the level sets of W . Part 2 follows by setting $g(t) = t^2$, $Z = R(M)$, and letting F be the σ -algebra generated by the level sets of W . \square

To construct an exchangeable pair to be used in our applications, we use the heat kernel on G . See [10], [25], [26], [16]) for a detailed discussion of heat kernels on compact Lie groups, including all of the properties stated in the remainder of this section. The papers [14], [1], [20], [15] illustrate combinatorial uses of heat kernels on compact Lie groups.

The heat kernel on G is defined by setting for $x, y \in G$ and $t \geq 0$,

$$(1) \quad K(t, x, y) = \sum_{n \geq 0} e^{-\lambda_n t} \phi_n(x) \overline{\phi_n(y)},$$

where the λ_n are the eigenvalues of the Laplacian repeated according to multiplicity, and the ϕ_n are an orthonormal basis of eigenfunctions of $L^2(G)$; these can be taken to be the irreducible characters of G .

We use the following properties of the heat kernel. Here Δ denotes the Laplacian of G , and $e^{t\Delta}$ is defined as $I + t\Delta + t^2 \frac{\Delta^2}{2!} + \dots$.

Lemma 2.3. *Let G be a compact Lie group, $x, y \in G$, and $t \geq 0$.*

(1) *$K(t, x, y)$ converges and is non-negative for all x, y, t .*

- (2) $\int_{y \in G} K(t, x, y) dy = 1$, where the integration is with respect to Haar measure of G .
- (3) $e^{t\Delta} \phi(x) = \int_{y \in G} K(t, x, y) \phi(y) dy$ for smooth ϕ .

The symmetry in x and y of $K(t, x, y)$ shows that the heat kernel is a reversible Markov process with respect to the Haar measure of G . Thus, given a function f on G , one can construct an exchangeable pair (W, W') by letting $W = f(M)$ where M is chosen from Haar measure, and $W' = f(M')$, where M' is obtained by moving time t from M via the heat kernel.

3. THE ORTHOGONAL GROUP

If λ is an integer partition (possibly with negative parts) and m_j denotes the multiplicity of part j in λ , we define $p_\lambda(M) = \prod_j \text{Tr}(M^j)^{m_j}$. For example, $p_{5,3,3}(M) = \text{Tr}(M^5) \text{Tr}(M^3)^2$. Typically we suppress the M and use the notation p_λ . We let $W = \frac{p_j}{\sqrt{j}}$ if j is odd and let $W = \frac{p_j - 1}{\sqrt{j}}$ if j is even. Note that since the eigenvalues of M are roots of unity and come in conjugate pairs, $p_j = p_{-j}$ is real. The main result of this section is a central limit theorem for W with error term $O(j/n)$.

The following moment computation of [11] (analogous to that of [5] for the full orthogonal group) will be helpful. In fact as the reader will see, in the applications of Lemma 3.1, we only use fourth moments and lower.

Lemma 3.1. *Let M be Haar distributed on $SO(n, \mathbb{R})$. Let (a_1, a_2, \dots, a_k) be a vector of non-negative integers. Let Z_1, \dots, Z_k be independent standard normal random variables. Let η_j be 1 if j is even and 0 otherwise. Then if $n - 1 \geq \sum_{i=1}^k a_i$,*

$$\mathbb{E} \left[\prod_{j=1}^k \text{Tr}(M^j)^{a_j} \right] = \prod_{j=1}^k g_j(a_j) = \prod_{j=1}^k \mathbb{E}(\sqrt{j} Z_j + \eta_j)^{a_j},$$

Here

$$\text{if } j \text{ is odd, } g_j(a) = \begin{cases} 0 & \text{if } a \text{ is odd} \\ j^{a/2} (a-1)(a-3) \cdots 1 & \text{if } a \text{ is even} \end{cases}$$

$$\text{if } j \text{ is even, } g_j(a) = 1 + \sum_{k \geq 1} \binom{a}{2k} j^k (2k-1)(2k-3) \cdots 1.$$

Rains [20] (see also [14]) determined how the Laplacian acts on power sum symmetric functions. We need his formula only in the following two cases.

Lemma 3.2. (1)

$$\Delta_{SO(n)} p_j = -\frac{(n-1)j}{2} p_j - \frac{j}{2} \sum_{1 \leq l < j} p_{l,j-l} + \frac{j}{2} \sum_{1 \leq l < j} p_{2l-j}.$$

(2)

$$\Delta_{SO(n)} p_{j,j} = -(n-1)j p_{j,j} - j^2 p_{2j} - j p_j \sum_{1 \leq l < j} p_{l,j-l} + j p_j \sum_{1 \leq l < j} p_{2l-j} + j^2 n.$$

We fix $t > 0$, and motivated by Section 2, define

$$W' = e^{t\Delta}(W) = W + \sum_{k \geq 1} \frac{t^k}{k!} \Delta^k(W).$$

Lemma 3.3 computes the conditional expectation $\mathbb{E}[W'|M]$.

Lemma 3.3.

$$\mathbb{E}[W'|M] = \left(1 - \frac{t(n-1)j}{2}\right) W + R(M),$$

with

$$R(M) = t \left[-\frac{\sqrt{j}}{2} \sum_{1 \leq l < j} p_{l,j-l} + \frac{\sqrt{j}}{2} \sum_{1 \leq l < j} p_{2l-j} \right] + O(t^2) \quad j \text{ odd},$$

and

$$R(M) = t \left[-\frac{(n-1)\sqrt{j}}{2} - \frac{\sqrt{j}}{2} \sum_{1 \leq l < j} p_{l,j-l} + \frac{\sqrt{j}}{2} \sum_{1 \leq l < j} p_{2l-j} \right] + O(t^2) \quad j \text{ even}.$$

Proof. Applying part 3 of Lemma 2.3 and part 1 of Lemma 3.2,

$$\begin{aligned} & \mathbb{E}[W'|M] \\ &= e^{t\Delta}(W) \\ &= W + t \left[-\frac{(n-1)\sqrt{j}}{2} p_j - \frac{\sqrt{j}}{2} \sum_{1 \leq l < j} p_{l,j-l} + \frac{\sqrt{j}}{2} \sum_{1 \leq l < j} p_{2l-j} \right] + O(t^2), \end{aligned}$$

and the result follows. \square

Lemma 3.4 computes $\mathbb{E}[(W' - W)^2|M]$, a quantity needed to apply Theorem 2.1. Many cancelations occur, and a simple formula emerges.

Lemma 3.4.

$$\mathbb{E}[(W' - W)^2|M] = tj(n - p_{2j}) + O(t^2).$$

Proof. Clearly

$$\mathbb{E}[(W' - W)^2|M] = \mathbb{E}[(W')^2|M] - 2W\mathbb{E}[W'|M] + W^2.$$

Suppose now that j is odd. By part 3 of Lemma 2.3 and part 2 of Lemma 3.2,

$$\begin{aligned} & \mathbb{E}[(W')^2|M] \\ &= W^2 + \frac{t}{j} \Delta p_{j,j} + O(t^2) \\ &= W^2 + t \left[-(n-1)p_{j,j} - jp_{2j} - p_j \sum_{1 \leq l < j} p_{l,j-l} + p_j \sum_{1 \leq l < j} p_{2l-j} + jn \right] \\ & \quad + O(t^2). \end{aligned}$$

By Lemma 3.3, $-2W\mathbb{E}[W'|M]$ is equal to

$$-2W^2 + t \left[(n-1)jW^2 + p_j \sum_{1 \leq l < j} p_{l,j-l} - p_j \sum_{1 \leq l < j} p_{2l-j} \right] + O(t^2).$$

Thus

$$\mathbb{E}[(W')^2|M] - 2W\mathbb{E}[W'|M] + W^2 = tj(n - p_{2j}) + O(t^2),$$

as claimed. A very similar calculation shows that the same conclusion holds for j even. \square

Lemma 3.5. *Suppose that $4j \leq n-1$. Then*

$$\text{Var}(\mathbb{E}[(W' - W)^2|M]) = 2j^3t^2 + O(t^3).$$

Proof. By Lemma 3.4,

$$\text{Var}(\mathbb{E}[(W' - W)^2|M]) = j^2t^2\text{Var}(p_{2j}) + O(t^3).$$

The result now follows from Lemma 3.1. \square

Lemma 3.6. *Suppose that $4j \leq n-1$. Then*

- (1) $\mathbb{E}(W' - W)^2 = tj(n-1) + O(t^2)$.
- (2) $\mathbb{E}(W' - W)^4 = O(t^2)$.

Proof. Lemma 3.4 implies that $\mathbb{E}(W' - W)^2 = \mathbb{E}[tj(n - p_{2j})] + O(t^2)$. From Lemma 3.1, $\mathbb{E}(p_{2j}) = 1$, which proves part 1 (only using that $2j \leq n-1$).

For part 2, first note that since

$\mathbb{E}[(W' - W)^4] = \mathbb{E}(W^4) - 4\mathbb{E}(W^3W') + 6\mathbb{E}[W^2(W')^2] - 4\mathbb{E}[W(W')^3] + \mathbb{E}[(W')^4]$,
exchangeability of (W, W') gives that

$$\begin{aligned} \mathbb{E}(W' - W)^4 &= 2\mathbb{E}(W^4) - 8\mathbb{E}(W^3W') + 6\mathbb{E}[W^2(W')^2] \\ &= 2\mathbb{E}(W^4) - 8\mathbb{E}[W^3\mathbb{E}[W'|M]] + 6\mathbb{E}[W^2\mathbb{E}[(W')^2|M]]. \end{aligned}$$

Supposing that j is odd and using Lemma 3.2, this simplifies to

$$\begin{aligned} &2\mathbb{E}(W^4) - 8\mathbb{E}[W^4] + 6\mathbb{E}[W^4] \\ &+ t\mathbb{E} \left[4(n-1)jW^4 + 4W^3\sqrt{j} \sum_{1 \leq l < j} p_{l,j-l} - 4W^3\sqrt{j} \sum_{1 \leq l < j} p_{2l-j} \right] \\ &+ t\mathbb{E} \left[-6(n-1)jW^4 - 6W^2p_j \sum_{1 \leq l < j} p_{l,j-l} + 6W^2p_j \sum_{1 \leq l < j} p_{2l-j} \right] \\ &+ t\mathbb{E} [-6jW^2p_{2j} + 6W^2jn] + O(t^2). \end{aligned}$$

By Lemma 3.1, this simplifies to

$$t [12j(n-1) - 18j(n-1) - 6j + 6jn] + O(t^2) = O(t^2),$$

as claimed. A very similar calculation gives the same conclusion for j even. \square

Next we bound a quantity appearing in the second term of Theorem 2.1.

Lemma 3.7. *Suppose that $4j \leq n - 1$. Let*

$$R(M) = t \left[-\frac{\sqrt{j}}{2} \sum_{1 \leq l < j} p_{l,j-l} + \frac{\sqrt{j}}{2} \sum_{1 \leq l < j} p_{2l-j} \right] + O(t^2) \quad j \text{ odd},$$

and

$$R(M) = t \left[-\frac{(n-1)\sqrt{j}}{2} - \frac{\sqrt{j}}{2} \sum_{1 \leq l < j} p_{l,j-l} + \frac{\sqrt{j}}{2} \sum_{1 \leq l < j} p_{2l-j} \right] + O(t^2) \quad j \text{ even}.$$

Then $\mathbb{E}[R^2] = O(t^2 j^4)$.

Proof. Suppose that j is odd. Applying Lemma 3.1 and keeping only terms with non-0 expectations, one has that

$$\begin{aligned} \mathbb{E}[R^2] &= \frac{t^2 j}{4} \mathbb{E} \left[4 \sum_{\substack{1 \leq l < j \\ l \text{ odd}}} (p_{l,j-l})^2 + 4 \sum_{\substack{1 \leq l < j \\ l \text{ odd}}} (p_l)^2 - 8 \sum_{\substack{1 \leq l < j \\ l \text{ odd}}} p_{l,j-l} \right] + O(t^3) \\ &= \frac{t^2 j}{4} \left[4 \sum_{\substack{1 \leq l < j \\ l \text{ odd}}} l(j-l+1) - 4 \sum_{\substack{1 \leq l < j \\ l \text{ odd}}} l \right] + O(t^3) \\ &= O(t^2 j^4). \end{aligned}$$

The case of j even is proved in a similar way, as can be seen by writing

$$R = t \left[\frac{\sqrt{j}}{2} - \frac{\sqrt{j}}{2} \sum_{1 \leq l < j} p_{l,j-l} + \frac{\sqrt{j}}{2} \sum_{\substack{1 \leq l < j \\ l \neq j/2}} p_{2l-j} \right] + O(t^2).$$

□

Combining the above calculations leads to the main result of this section.

Theorem 3.8. *Let M be chosen from the Haar measure of $SO(n, \mathbb{R})$. Let $W(M) = \frac{\text{Tr}(M^j)}{\sqrt{j}}$ if j is odd and $W(M) = \frac{\text{Tr}(M^j)-1}{\sqrt{j}}$ if j is even. Then*

$$\left| \mathbb{P}(W \leq x_0) - \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{x_0} e^{-\frac{x^2}{2}} dx \right| = O(j/n).$$

Proof. The result is trivial if $4j > n - 1$, so assume that $4j \leq n - 1$. We apply Theorem 2.1 to the exchangeable pair (W, W') with $a = \frac{t(n-1)j}{2}$, and will take the limit $t \rightarrow 0$ in each term (keeping j, n fixed). By part 1 of Lemma 2.2 and Lemma 3.5, the first term is $O(\sqrt{j}/n)$. By part 2 of Lemma 2.2 and Lemma 3.7, the second term is $O(j/n)$. By the Cauchy-Schwarz inequality and Lemma 3.6,

$$\mathbb{E}|W' - W|^3 \leq \sqrt{\mathbb{E}(W' - W)^2 \mathbb{E}(W' - W)^4} = O(t^{3/2}).$$

Thus the third term in Theorem 2.1 tends to 0 as $t \rightarrow 0$, and the result is proved. □

4. THE SYMPLECTIC GROUP

Let J be the $2n \times 2n$ matrix of the form $\begin{pmatrix} 0 & I \\ -I & 0 \end{pmatrix}$ with all blocks $n \times n$. $USp(2n, \mathbb{C})$ is defined as the set of $2n \times 2n$ unitary matrices M with complex entries such that $MJM^t = J$; it consists of the matrices preserving an alternating form. As in Section 3, we use the notation that $p_\lambda(M) = \prod_j \text{Tr}(M^j)^{m_j}$, and we typically suppress the M and use the notation p_λ . We let $W = \frac{p_j}{\sqrt{j}}$ if j is odd and let $W = \frac{p_{j+1}}{\sqrt{j}}$ if j is even. Since the eigenvalues of M are roots of unity and come in conjugate pairs, $p_j = p_{-j}$ is real valued. The main result of this section is a central limit theorem for W with error term $O(j/n)$.

The following moment computation is the symplectic analog of Lemma 3.1. It was proved by [5] under the slightly weaker assumption that $n \geq \sum_{i=1}^k a_i$. As stated, Lemma 4.1 appears in [11], with a later proof in [19].

Lemma 4.1. *Let M be Haar distributed on $USp(2n, \mathbb{C})$. Let (a_1, a_2, \dots, a_k) be a vector of non-negative integers. Let Z_1, \dots, Z_k be independent standard normal random variables. Let η_j be 1 if j is even and 0 otherwise. Then if $2n + 1 \geq \sum_{i=1}^k a_i$,*

$$\mathbb{E} \left[\prod_{j=1}^k \text{Tr}(M^j)^{a_j} \right] = \prod_{j=1}^k (-1)^{(j-1)a_j} g_j(a_j) = \prod_{j=1}^k \mathbb{E}(\sqrt{j}Z_j - \eta_j)^{a_j},$$

where the polynomials g_j are as in Lemma 3.1.

Rains [20] (see also [14]) determined how the Laplacian acts on power sum symmetric functions. We need his formula only in the following two cases.

Lemma 4.2. (1)

$$\Delta_{USp(2n)} p_j = -\frac{(2n+1)j}{2} p_j - \frac{j}{2} \sum_{1 \leq l < j} p_{2l-j} - \frac{j}{2} \sum_{1 \leq l < j} p_{l,j-l}.$$

(2)

$$\Delta_{USp(2n)} p_{j,j} = -(2n+1)j p_{j,j} - j^2 p_{2j} - j p_j \sum_{1 \leq l < j} p_{2l-j} - j p_j \sum_{1 \leq l < j} p_{l,j-l} + 2j^2 n.$$

As in the orthogonal case, we fix $t > 0$, and define

$$W' = e^{t\Delta}(W) = W + \sum_{k \geq 1} \frac{t^k}{k!} \Delta^k(W).$$

Lemma 4.3.

$$\mathbb{E}[W'|M] = \left(1 - \frac{t(2n+1)j}{2}\right) W + R(M),$$

with

$$R(M) = t \left[-\frac{\sqrt{j}}{2} \sum_{1 \leq l < j} p_{2l-j} - \frac{\sqrt{j}}{2} \sum_{1 \leq l < j} p_{l,j-l} \right] + O(t^2) \quad j \text{ odd},$$

and

$$R(M) = t \left[\frac{(2n+1)\sqrt{j}}{2} - \frac{\sqrt{j}}{2} \sum_{1 \leq l < j} p_{2l-j} - \frac{\sqrt{j}}{2} \sum_{1 \leq l < j} p_{l,j-l} \right] + O(t^2) \quad j \text{ even.}$$

Proof. Applying part 3 of Lemma 2.3 and part 1 of Lemma 4.2,

$$\begin{aligned} & \mathbb{E}[W'|W] \\ &= e^{t\Delta}(W) \\ &= W + t \left[-\frac{(2n+1)\sqrt{j}}{2} p_j - \frac{\sqrt{j}}{2} \sum_{1 \leq l < j} p_{2l-j} - \frac{\sqrt{j}}{2} \sum_{1 \leq l < j} p_{l,j-l} \right] + O(t^2), \end{aligned}$$

and the result follows. \square

Lemma 4.4 computes $\mathbb{E}[(W' - W)^2|M]$, a quantity needed to apply Theorem 2.1. As in the orthogonal case, there are many cancelations, leading to a simple formula.

Lemma 4.4.

$$\mathbb{E}[(W' - W)^2|M] = tj(2n - p_{2j}) + O(t^2).$$

Proof. Clearly

$$\mathbb{E}[(W' - W)^2|M] = \mathbb{E}[(W')^2|M] - 2W\mathbb{E}[W'|M] + W^2.$$

Suppose that j is odd. By part 3 of Lemma 2.3 and part 2 of Lemma 4.2,

$$\begin{aligned} & \mathbb{E}[(W')^2|M] \\ &= W^2 + \frac{t}{j} \Delta p_{j,j} + O(t^2) \\ &= W^2 + t \left[-(2n+1)p_{j,j} - jp_{2j} - p_j \sum_{1 \leq l < j} p_{2l-j} - p_j \sum_{1 \leq l < j} p_{l,j-l} + 2jn \right] \\ & \quad + O(t^2). \end{aligned}$$

By Lemma 4.3, $-2W\mathbb{E}[W'|M]$ is equal to

$$-2W^2 + t \left[(2n+1)p_{j,j} + p_j \sum_{1 \leq l < j} p_{2l-j} + p_j \sum_{1 \leq l < j} p_{l,j-l} \right] + O(t^2).$$

Thus

$$\mathbb{E}[(W')^2|M] - 2W\mathbb{E}[W'|M] + W^2 = tj[2n - p_{2j}] + O(t^2),$$

as needed. A similar computation proves the lemma for j even. \square

Lemma 4.5. *Suppose that $4j \leq 2n+1$. Then*

$$\text{Var}(\mathbb{E}[(W' - W)^2|M]) = 2j^3t^2 + O(t^3).$$

Proof. By Lemma 4.4,

$$\text{Var}(\mathbb{E}[(W' - W)^2|M]) = j^2t^2 \text{Var}(p_{2j}) + O(t^3).$$

The result now follows from Lemma 4.1. \square

Lemma 4.6. *Suppose that $4j \leq 2n + 1$.*

- (1) $\mathbb{E}(W' - W)^2 = tj(2n + 1) + O(t^2)$.
- (2) $\mathbb{E}(W' - W)^4 = O(t^2)$.

Proof. Lemma 4.4 implies that $\mathbb{E}(W' - W)^2 = \mathbb{E}[tj(2n - p_{2j})] + O(t^2)$. From Lemma 4.1, $\mathbb{E}(p_{2j}) = -1$, which proves part 1 (even assuming that $2j \leq 2n + 1$).

For part 2, first note that since

$$\mathbb{E}[(W' - W)^4] = \mathbb{E}(W^4) - 4\mathbb{E}(W^3W') + 6\mathbb{E}[W^2(W')^2] - 4\mathbb{E}[W(W')^3] + \mathbb{E}[(W')^4],$$

exchangeability of (W, W') gives that

$$\begin{aligned} \mathbb{E}(W' - W)^4 &= 2\mathbb{E}(W^4) - 8\mathbb{E}(W^3W') + 6\mathbb{E}[W^2(W')^2] \\ &= 2\mathbb{E}(W^4) - 8\mathbb{E}[W^3\mathbb{E}[W'|M]] + 6\mathbb{E}[W^2\mathbb{E}[(W')^2|M]]. \end{aligned}$$

Suppose j is odd. Using Lemma 4.3 and part 2 of Lemma 4.2, this becomes

$$\begin{aligned} &2\mathbb{E}(W^4) - 8\mathbb{E}(W^4) + 6\mathbb{E}(W^4) \\ &+ t\mathbb{E} \left[4(2n + 1)jW^4 + 4W^3\sqrt{j} \sum_{1 \leq l < j} p_{l,j-l} + 4W^3\sqrt{j} \sum_{1 \leq l < j} p_{2l-j} \right] \\ &+ t\mathbb{E} \left[-6(2n + 1)jW^4 - 6jW^2p_{2j} - 6W^2p_j \sum_{1 \leq l < j} p_{l,j-l} \right] \\ &+ t \left[-6W^2p_j \sum_{1 \leq l < j} p_{2l-j} + 12W^2jn \right] + O(t^2). \end{aligned}$$

By Lemma 4.1, this simplifies to

$$t[12j(2n + 1) - 18j(2n + 1) + 6j + 12jn] + O(t^2) = O(t^2),$$

as claimed. A similar calculation gives the same result for j even. \square

Lemma 4.7. *Suppose that $4j \leq 2n + 1$. Let*

$$R(M) = t \left[-\frac{\sqrt{j}}{2} \sum_{1 \leq l < j} p_{2l-j} - \frac{\sqrt{j}}{2} \sum_{1 \leq l < j} p_{l,j-l} \right] + O(t^2) \quad j \text{ odd},$$

and

$$R(M) = t \left[\frac{(2n + 1)\sqrt{j}}{2} - \frac{\sqrt{j}}{2} \sum_{1 \leq l < j} p_{2l-j} - \frac{\sqrt{j}}{2} \sum_{1 \leq l < j} p_{l,j-l} \right] + O(t^2) \quad j \text{ even}.$$

Then $\mathbb{E}[R^2] = O(t^2j^4)$.

Proof. Suppose that j is odd. Applying Lemma 4.1 and keeping only terms with non-0 contribution, one has that

$$\begin{aligned}\mathbb{E}[R^2] &= \frac{t^2 j}{4} \mathbb{E} \left[4 \sum_{\substack{1 \leq l < j \\ l \text{ odd}}} (p_{l,j-l})^2 + 4 \sum_{\substack{1 \leq l < j \\ l \text{ odd}}} (p_l)^2 + 8 \sum_{\substack{1 \leq l < j \\ l \text{ odd}}} p_{l,j-l} \right] + O(t^3) \\ &= \frac{t^2 j}{4} \left[4 \sum_{\substack{1 \leq l < j \\ l \text{ odd}}} l(j-l+1) - 4 \sum_{\substack{1 \leq l < j \\ l \text{ odd}}} l \right] + O(t^3) \\ &= O(t^2 j^4).\end{aligned}$$

The case of j even is proved by a similar argument, after writing

$$R = t \left[\frac{\sqrt{j}}{2} - \frac{\sqrt{j}}{2} \sum_{\substack{1 \leq l < j \\ l \neq j/2}} p_{2l-j} - \frac{\sqrt{j}}{2} \sum_{1 \leq l < j} p_{l,j-l} \right] + O(t^2).$$

□

Theorem 4.8. *Let M be chosen from the Haar measure of $USp(2n, \mathbb{C})$. Let $W(M) = \frac{Tr(M^j)}{\sqrt{j}}$ if j is odd, and $W(M) = \frac{Tr(M^j)+1}{\sqrt{j}}$ if j is even. Then*

$$\left| \mathbb{P}(W \leq x_0) - \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{x_0} e^{-\frac{x^2}{2}} dx \right| = O(j/n).$$

Proof. The result is trivial if $4j > 2n + 1$, so assume that $4j \leq 2n + 1$. We apply Theorem 2.1 to the exchangeable pair (W, W') with $a = \frac{t(2n+1)j}{2}$, and will take the limit $t \rightarrow 0$ in each term (keeping j, n fixed). By part 1 of Lemma 2.2 and Lemma 4.5, the first term is $O(\sqrt{j}/n)$. By part 2 of Lemma 2.2 and Lemma 4.7, the second term is $O(j/n)$. By the Cauchy-Schwarz inequality and Lemma 4.6,

$$\mathbb{E}|W' - W|^3 \leq \sqrt{\mathbb{E}(W' - W)^2 \mathbb{E}(W' - W)^4} = O(t^{3/2}).$$

Thus the third term in Theorem 2.1 tends to 0 as $t \rightarrow 0$, and the result follows. □

5. THE UNITARY GROUP

In this final section, we treat the unitary group $U(n, \mathbb{C})$. We let p_λ be as in Section 3 and Section 4 and define the real valued random variable $W = \frac{p_j + \bar{p}_j}{\sqrt{2j}}$. The main result of this section is a central limit theorem for W , with error term $O(j/n)$. To begin, we recall the following moment computation from [5].

Lemma 5.1. *Let M be Haar distributed on $U(n, \mathbb{C})$. Let (a_1, a_2, \dots, a_k) and (b_1, \dots, b_k) be vectors of non-negative integers. Let Z_1, \dots, Z_k be independent standard normal random variables. Then for all $n \geq \sum_{i=1}^k (a_i + b_i)$,*

$$\mathbb{E} \left[\prod_{j=1}^k Tr(M^j)^{a_j} \cdot \overline{Tr(M^j)^{b_j}} \right] = \delta_{\vec{a}\vec{b}} \prod_{j=1}^k j^{a_j} a_j!.$$

Rains [20] (see also [14]) determined how the Laplacian acts on power sum symmetric functions. We require his formulas only in the following cases.

Lemma 5.2. (1)

$$\Delta_{U(n)} p_j = -nj p_j - j \sum_{1 \leq l < j} p_{l,j-l}.$$

(2)

$$\Delta_{U(n)} p_{j,j} = -2nj p_{j,j} - 2j^2 p_{2j} - 2j p_j \sum_{1 \leq l < j} p_{l,j-l}.$$

(3)

$$\Delta_{U(n)} (p_j \overline{p_j}) = 2j^2 n - 2nj p_j \overline{p_j} - j p_j \sum_{1 \leq l < j} \overline{p_{l,j-l}} - j \overline{p_j} \sum_{1 \leq l < j} p_{l,j-l}.$$

Lemma 5.3 computes the conditional expectation $\mathbb{E}[W'|M]$.

Lemma 5.3.

$$\mathbb{E}[W'|M] = (1 - njt) W + R(M),$$

with

$$R(M) = t \left[-\sqrt{\frac{j}{2}} \sum_{1 \leq l < j} p_{l,j-l} - \sqrt{\frac{j}{2}} \sum_{1 \leq l < j} \overline{p_{l,j-l}} \right] + O(t^2).$$

Proof. Applying Lemma 2.3 and part 1 of Lemma 5.2 gives that

$$\begin{aligned} & \mathbb{E}[W'|M] \\ &= e^{t\Delta}(W) \\ &= W + t \left[-njW - \sqrt{\frac{j}{2}} \sum_{1 \leq l < j} p_{l,j-l} - \sqrt{\frac{j}{2}} \sum_{1 \leq l < j} \overline{p_{l,j-l}} \right] + O(t^2), \end{aligned}$$

as desired. \square

Lemma 5.4 computes $\mathbb{E}[(W' - W)^2|M]$. As in the other cases, there are nice cancelations.

Lemma 5.4.

$$\mathbb{E}[(W' - W)^2|M] = tj(2n - p_{2j} - \overline{p_{2j}}) + O(t^2).$$

Proof. Clearly

$$\mathbb{E}[(W' - W)^2|M] = \mathbb{E}[(W')^2|M] - 2W\mathbb{E}[W'|M] + W^2.$$

By Lemmas 2.3 and 5.2,

$$\begin{aligned}
& \mathbb{E}[(W')^2|M] \\
&= W^2 + \frac{t}{2j} \Delta[p_{j,j} + 2p_j \overline{p_j} + \overline{p_{j,j}}] + O(t^2) \\
&= W^2 + t \left[-np_{j,j} - jp_{2j} - p_j \sum_{1 \leq l < j} p_{l,j-l} - n\overline{p_{j,j}} - j\overline{p_{2j}} \right] \\
&\quad + t \left[-\overline{p_j} \sum_{1 \leq l < j} \overline{p_{l,j-l}} + 2jn - 2np_j \overline{p_j} - p_j \sum_{1 \leq l < j} \overline{p_{l,j-l}} - \overline{p_j} \sum_{1 \leq l < j} p_{l,j-l} \right] \\
&\quad + O(t^2).
\end{aligned}$$

By Lemma 5.3, $-2W\mathbb{E}[W'|M]$ is equal to

$$\begin{aligned}
& -2W^2 + t \left[np_{j,j} + 2np_j \overline{p_j} + n\overline{p_{j,j}} + p_j \sum_{1 \leq l < j} p_{l,j-l} \right] \\
& + t \left[p_j \sum_{1 \leq l < j} \overline{p_{l,j-l}} + \overline{p_j} \sum_{1 \leq l < j} p_{l,j-l} + \overline{p_j} \sum_{1 \leq l < j} \overline{p_{l,j-l}} \right] + O(t^2).
\end{aligned}$$

Thus

$$\mathbb{E}[(W')^2|M] - 2W\mathbb{E}[W'|M] + W^2 = tj[2n - p_{2j} - \overline{p_{2j}}] + O(t^2),$$

and the lemma is proved. \square

Lemma 5.5. *Suppose that $4j \leq n$. Then*

$$\text{Var}(\mathbb{E}[(W' - W)^2|M]) = 4j^3 t^2 + O(t^3).$$

Proof. By Lemmas 5.4 and 5.1,

$$\begin{aligned}
\text{Var}(\mathbb{E}[(W' - W)^2|M]) &= j^2 t^2 \text{Var}(p_{2j} + \overline{p_{2j}}) + O(t^3) \\
&= j^2 t^2 \mathbb{E}[(p_{2j} + \overline{p_{2j}})^2] + O(t^3) \\
&= 4j^3 t^2 + O(t^3).
\end{aligned}$$

\square

Lemma 5.6. *Suppose that $4j \leq n$.*

- (1) $\mathbb{E}(W' - W)^2 = t2jn + O(t^2)$.
- (2) $\mathbb{E}(W' - W)^4 = O(t^2)$.

Proof. Lemma 5.4 implies that $\mathbb{E}(W' - W)^2 = \mathbb{E}[tj(2n - p_{2j} - \overline{p_{2j}})] + O(t^2)$. From Lemma 5.1, $\mathbb{E}(p_{2j}) = \mathbb{E}(\overline{p_{2j}}) = 0$, which proves part 1 (using only that $2j \leq n$).

For part 2, first note that since

$$\mathbb{E}[(W' - W)^4] = \mathbb{E}(W^4) - 4\mathbb{E}(W^3 W') + 6\mathbb{E}[W^2 (W')^2] - 4\mathbb{E}[W (W')^3] + \mathbb{E}[(W')^4],$$

exchangeability of (W, W') gives that

$$\begin{aligned}
\mathbb{E}(W' - W)^4 &= 2\mathbb{E}(W^4) - 8\mathbb{E}(W^3 W') + 6\mathbb{E}[W^2 (W')^2] \\
&= 2\mathbb{E}(W^4) - 8\mathbb{E}[W^3 \mathbb{E}[W'|M]] + 6\mathbb{E}[W^2 \mathbb{E}[(W')^2|M]].
\end{aligned}$$

Using Lemmas 5.2 and 5.3, this simplifies to

$$\begin{aligned}
& 2\mathbb{E}[W^4] - 8\mathbb{E}[W^4] + 6\mathbb{E}[W^4] \\
& + t\mathbb{E} \left[8njW^4 + 8W^3 \sqrt{\frac{j}{2}} \sum_{1 \leq l < j} p_{l,j-l} + 8W^3 \sqrt{\frac{j}{2}} \sum_{1 \leq l < j} \overline{p_{l,j-l}} \right] \\
& + t\mathbb{E} \left[-6nW^2 p_{j,j} - 6jW^2 p_{2j} - 6W^2 p_j \sum_{1 \leq l < j} p_{l,j-l} - 6nW^2 \overline{p_{j,j}} \right] \\
& + t\mathbb{E} \left[-6jW^2 \overline{p_{2j}} - 6W^2 \overline{p_j} \sum_{1 \leq l < j} \overline{p_{l,j-l}} + 12njW^2 \right] \\
& + t\mathbb{E} \left[-12nW^2 p_j \overline{p_j} - 6W^2 p_j \sum_{1 \leq l < j} \overline{p_{l,j-l}} - 6W^2 \overline{p_j} \sum_{1 \leq l < j} p_{l,j-l} \right] + O(t^2).
\end{aligned}$$

By Lemma 5.1, after dropping out terms with 0 expectation, there remains

$$\begin{aligned}
& t\mathbb{E}[8W^4 jn - 6W^2 n p_{j,j} - 6W^2 n \overline{p_{j,j}} + 12W^2 jn - 12W^2 n p_j \overline{p_j}] + O(t^2) \\
& = t[24jn - 6jn - 6jn + 12jn - 24jn] + O(t^2) \\
& = O(t^2),
\end{aligned}$$

as needed. \square

Lemma 5.7. *Let $R = t \left[-\sqrt{\frac{j}{2}} \sum_{1 \leq l < j} p_{l,j-l} - \sqrt{\frac{j}{2}} \sum_{1 \leq l < j} \overline{p_{l,j-l}} \right] + O(t^2)$, and suppose that $4j \leq n$. Then $\mathbb{E}[R^2] \leq \frac{j^4 t^2}{4} + O(t^3)$.*

Proof. Applying Lemma 5.1 and keeping only terms with non-0 contribution, one has that

$$\mathbb{E}[R^2] = jt^2 \mathbb{E} \left[\sum_{1 \leq l < j} p_{l,j-l} \overline{p_{l,j-l}} \right] + O(t^3).$$

If j is odd, then by Lemma 5.1,

$$\mathbb{E}[R^2] = jt^2 \sum_{1 \leq l < j} [l(j-l)] + O(t^3) = \frac{(j^4 - j^2)}{6} t^2 + O(t^3),$$

while if j is even, one obtains that

$$\mathbb{E}[R^2] = \frac{(2j^4 + 3j^3 - 2j^2)}{12} t^2 + O(t^3).$$

The result follows. \square

Theorem 5.8. *Let M be chosen from the Haar measure of $U(n, \mathbb{C})$, and let $W(M) = \frac{1}{\sqrt{2j}} [Tr(M^j) + \overline{Tr(M^j)}]$. Then*

$$\left| \mathbb{P}(W \leq x_0) - \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{x_0} e^{-\frac{x^2}{2}} dx \right| = O(j/n).$$

Proof. The result is trivial if $4j > n$, so assume that $4j \leq n$. We apply Theorem 2.1 to the exchangeable pair (W, W') with $a = tnj$, and will take the limit $t \rightarrow 0$ in each term. By part 1 of Lemma 2.2 and Lemma 5.5, the first term is at most $\frac{12\sqrt{j}}{n}$. By Lemma 5.7 and part 2 of Lemma 2.2, the second term in Theorem 2.1 is at most $\frac{19j}{2n}$. By the Cauchy-Schwarz inequality and Lemma 5.6,

$$\mathbb{E}|W' - W|^3 \leq \sqrt{\mathbb{E}(W' - W)^2 \mathbb{E}(W' - W)^4} = O(t^{3/2}).$$

Thus the third term in Theorem 2.1 tends to 0 as $t \rightarrow 0$, and the result follows since

$$\frac{12\sqrt{j}}{n} + \frac{19j}{2n} \leq \frac{22j}{n}.$$

□

ACKNOWLEDGEMENTS

We thank Eric Rains for helpful correspondence. The author was partially supported by NSF grant DMS 0802082 and NSA grant H98230-08-1-0133.

REFERENCES

- [1] Biane, P., Free Brownian motion, free stochastic calculus and random matrices, in *Free probability theory (Waterloo, ON, 1995)*, 1-19, Fields Inst. Commun., 12, Amer. Math. Soc., Providence, RI, 1997.
- [2] Chatterjee, S., Fluctuations of eigenvalues and second order Poincaré inequalities, *Probab. Theory Related Fields* **143** (2009), 1-40.
- [3] Collins, B. and Stolz, M., Borel theorems for random matrices from the classical compact symmetric spaces, *Ann. Probab.* **36** (2008), 876-895.
- [4] Diaconis, P. and Evans, S., Linear functionals of eigenvalues of random matrices, *Trans. Amer. Math. Soc.* **353** (2001), 2615-2633 (electronic).
- [5] Diaconis, P. and Shahshahani, M., On the eigenvalues of random matrices. Studies in applied probability. *J. Appl. Probab.* **31A** (1994), 49-62.
- [6] Duits, M. and Johansson, K., Powers of large random unitary matrices and Toeplitz determinants, *Trans. Amer. Math. Soc.* **362** (2010), 1169-1187.
- [7] Dumitriu, I. and Edelman, A., Global spectrum fluctuations for the beta-Hermite and beta-Laguerre ensembles via matrix models, *J. Math. Phys.* **47** (2006) no. 6, 063302, 36 pp.
- [8] Durrett, R., Probability: theory and examples. Second edition. Duxbury Press, Belmont, CA, 1996.
- [9] Fulman, J., Stein's method and characters of compact Lie groups, *Comm. Math. Phys.* **288** (2009), 1181-1201.
- [10] Grigor'yan, A., Heat kernel and analysis on manifolds, AMS/IP Studies in Advanced Mathematics, 47. American Mathematical Society, Providence, RI; International Press, Boston, MA, 2009.
- [11] Hughes, C. P. and Rudnick, Z., Mock-Gaussian behaviour for linear statistics of classical compact groups. *Random matrix theory. J. Phys. A* **36** (2003), 2919-2932.

- [12] Johansson, K., On random matrices from the compact classical groups, *Ann. of Math.* **145** (1997), 519-545.
- [13] Jorgenson, J. and Lang, S., The ubiquitous heat kernel. Mathematics unlimited—2001 and beyond, 655–683, Springer, Berlin, 2001.
- [14] Levy, T., Schur-Weyl duality and the heat kernel measure on the unitary group, *Adv. Math.* **218** (2008), 537-575.
- [15] Liu, K., Heat kernels, symplectic geometry, moduli spaces and finite groups, in *Surveys in differential geometry: differential geometry inspired by string theory*, 527-542, Surv. Differ. Geom., 5, Int. Press, Boston, MA, 1999.
- [16] Maher, D., Brownian motion and heat kernels on compact Lie groups and symmetric spaces, Ph.D. thesis, University of New South Wales, 2006.
- [17] Meckes, E., On the approximate normality of eigenfunctions of the Laplacian, *Trans. Amer. Math. Soc.* **361** (2009), 5377-5399.
- [18] Meckes, E., An infinitesimal version of Stein's method of exchangeable pairs, Stanford University Ph.D. thesis, 2006.
- [19] Pastur, L. and Vasilchuk, V., On the moments of traces of matrices of classical groups, *Comm. Math. Phys.* **252** (2004), 149-166.
- [20] Rains, E. M., Combinatorial properties of Brownian motion on the compact classical groups, *J. Theoret. Probab.* **10** (1997), 659-679.
- [21] Rains, E. M., High powers of random elements of compact Lie groups, *Probab. Theory Related Fields* **107** (1997), 219-241.
- [22] Reinert, G., Couplings for normal approximations with Stein's method, in *Microsurveys in discrete probability (Princeton, NJ, 1997)*, 193-207, DIMACS Ser. Discrete Math. Theoret. Comput. Sci., 41, Amer. Math. Soc., Providence, RI, 1998.
- [23] Rinott, Y. and Rotar, V., Normal approximations by Stein's method, *Decis. Econ. Finance* **23** (2000), 15-29.
- [24] Rinott, Y. and Rotar, V., On coupling constructions and rates in the CLT for dependent summands with applications to the antivoter model and weighted U -statistics, *Ann. Appl. Probab.* **7** (1997), 1080-1105.
- [25] Rosenberg, S., The Laplacian on a Riemannian manifold. An introduction to analysis on manifolds. London Mathematical Society Student Texts, 31. Cambridge University Press, Cambridge, 1997.
- [26] Saloff-Coste, L., Precise estimates on the rate at which certain diffusions tend to equilibrium, *Math. Z.* **217** (1994), 641-677.
- [27] Sarnak, P., Arithmetic quantum chaos. *The Schur lectures* (1992) (Tel Aviv), Israel Math. Conf. Proc., 8, Bar-Ilan Univ., Ramat Gan, 1995.
- [28] Sinai, Y. and Soshnikov, A., Central limit theorem for traces of large random symmetric matrices with independent matrix elements, *Bol. Soc. Brasil. Mat. (N.S.)* **29** (1998), 1-24.
- [29] Soshnikov, A., The central limit theorem for local linear statistics in classical compact groups and related combinatorial identities, *Ann. Probab.* **28** (2000), 1353-1370.
- [30] Stein, C., The accuracy of the normal approximation to the distribution of the traces of powers of random orthogonal matrices. Stanford

- University Statistics Department technical report no. 470, (1995).
- [31] Stein, C., Approximate computation of expectations. Institute of Mathematical Statistics Lecture Notes-Monograph Series, 7. Institute of Mathematical Statistics, Hayward, CA, 1986.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF SOUTHERN CALIFORNIA, LOS ANGELES, CA, 90089, USA

E-mail address: `fulman@usc.edu`